

Chorus type whistler wave distributions in the radiation belts and inner magnetosphere using ray tracing

H. Breuillard¹, Y.Zaliznyak^{1,2}, O.Agapitov^{1,3}, V. Krasnoselskikh¹, A. Artemyev^{1,4}, G. Rolland⁵

¹LPC2E/CNRS University of Orleans, Orleans, France

²Institute for Nuclear Research, Kiev, Ukraine

³National Taras Shevchenko University of Kiev, Kiev, Ukraine

⁴Space Research Institute, RAS, Moscow, Russia

⁵CNES, Toulouse, France

mél: hugo.breuillard@cnr-orleans.fr

The quasi-monochromatic whistler wave packets are widely assumed to be formed in the vicinity of the magnetic equator and are frequently observed for example by Cluster spacecraft. The objective of our study is a reconstruction of realistic chorus emissions in radiation belts and in inner magnetosphere. To achieve this aim the data from the electric and magnetic field measurements onboard Cluster satellite [1] is used to determine the major characteristics of the chorus signal around the equator region, namely, its averaged wave vector, wave vector distribution, Poynting flux and polarization. Then the propagation of such a wave packet is modeled in the framework of so-called ray tracing technique using the original code which employs K. Rönmark's WHAMP to obtain hot plasma dispersion function values along the wave packet trajectory. The observed (real) rays at the equator are first fitted to the observed waveform using Cluster observations (initial conditions) and then these rays are propagated numerically through the inner magnetosphere in the frame of the WKB approximation. The density distributions of the magnetospheric particles is taken from the Gallagher et al. package that is provided by the authors and distributed as free software. Ray tracing allows one to reconstruct the properties of waves such as electric and magnetic fields, and the width of the wave packet in k-space along the ray propagation path. The calculations take into account realistic effects of the spreading of the signal due to propagation in the inhomogeneous and anisotropic magnetized plasma, the dependence of signal propagation characteristics upon initial conditions, etc. The obtained results are in a good agreement with the experimental wave normal dependence on latitude. Our calculations make possible to follow the wave packets and calculate their properties in the desired regions, e.g. the regions where an efficient wave-particle interaction is expected to occur (Figure 1). This allows to estimate particles diffusion rates [2].




Figure 1: PDF of $\tan\theta$ distribution from CLUSTER experimental data and from simulation.

Références

[1] Agapitov, O., V. Krasnoselskikh, Y. V. Khotyaintsev, and G. Rolland. GRL,38 (2011)

[2] Artemyev, A., O. Agapitov, H. Breuillard, V. Krasnoselskikh, G. Rolland GRL,40 (2012)